



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 908 453 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
**26.06.2002 Bulletin 2002/26**

(51) Int Cl.7: **C07D 257/04, C12Q 1/32,  
C07C 309/46**

(21) Application number: **97917413.3**

(86) International application number:  
**PCT/JP97/01324**

(22) Date of filing: **17.04.1997**

(87) International publication number:  
**WO 97/38985 (23.10.1997 Gazette 1997/45)**

### (54) **WATER-SOLUBLE TETRAZOLIUM SALT COMPOUNDS**

**WASSERLÖSLICHE TETRAZOLIUMSALZ-VERBINDUNGEN**

**COMPOSES DE SELS DE TETRAZOLIUM SOLUBLES DANS L'EAU**

(84) Designated Contracting States:  
**BE CH DE FR GB IT LI**

(30) Priority: **18.04.1996 JP 12113496**

(43) Date of publication of application:  
**14.04.1999 Bulletin 1999/15**

(73) Proprietor: **Dojindo Laboratories Co., Ltd.**  
**Kamimashiki-gun, Kumamoto 861-2202 (JP)**

(72) Inventors:  
• **ISHIYAMA, Munetaka**  
**Kumamoto 862 (JP)**  
• **MIYAZONO, Yoko**  
**Kumamoto 861-11 (JP)**  
• **SHIGA, Masanobu**  
**Kumamoto 862 (JP)**

• **SASAMOTO, Kazumi**  
**Kumamoto 861-22 (JP)**

(74) Representative: **Godemeyer, Thomas, Dr.**  
**Sternagel, Fleischer, Godemeyer & Partner**  
**Patentanwälte**  
**An den Gärten 7**  
**51491 Overath (DE)**

(56) References cited:  
**JP-A- 7 070 092** **JP-A- 56 061 366**  
**JP-A- 56 061 367**

• **CHEMICAL & PHARMACEUTICAL BULLETIN,**  
**Vol. 41, No. 6, (1993), ISHIYAMA M. et al., "A New**  
**Tetrazorium Salt that Produces a Highly**  
**Water-Soluble Formazan Dye", p. 1118-1122.**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 908 453 B1**




of low storage stability when stored in the state of an aqueous solution. Because aqueous solutions are often stored, after their preparation, for a long period of time such as for 3 to 6 months in usual clinical analyses, it is desired that tetrazolium compounds used as hydrogen acceptors have high storage stability in the state of an aqueous solution.

# 5 Disclosure of the Invention

[0009] The inventors of the present invention conducted various studies to provide compounds which have high storage stability as an aqueous solution, and in addition, maintains the characteristic excellent detection sensitivity of the compounds of the general formula (2). As a result, they found that the compounds of the general formula (1) set out below have the desired features. The present invention was achieved on the basis of these findings.

[0010] The present invention thus provides tetrazolium salt compounds represented by the following general formula (1):




wherein R<sup>1</sup> is a C<sub>1-4</sub> alkyl group or a C<sub>1-4</sub> alkoxy group and M represents an alkali metal or an ammonium.

[0011] According to preferred embodiments of the present invention, there are provided the aforementioned tetrazolium salt compounds wherein R<sup>1</sup> is methyl group or methoxy group, and M is sodium.

[0012] According to another aspect of the present invention, there is provided a reagent for the measurement of a dehydrogenase which comprises the aforementioned tetrazolium salt compound. As a preferred embodiment of the present invention, there is provided the aforementioned reagent which is used for a measurement utilizing reduced nicotinamide adenine dinucleotide as an intermediate electron transporter.

[0013] According to further aspect of the present invention, there are provided formazan compounds represented by the following general formula (6):



wherein R<sup>1</sup> is a C<sub>1-4</sub> alkyl group or a C<sub>1-4</sub> alkoxy group.

[0014] According to still further aspect of the present invention, there is provided a method for the measurement of a dehydrogenase wherein the compound of the above general formula (1) is used as a hydrogen acceptor. According to preferred embodiments of the aforementioned methods, there are provided the method which comprises the step of measuring the absorbance of the formazan compound represented by the above general formula (6); and the method wherein reduced nicotinamide adenine dinucleotide is used as an intermediate electronic transporter.

## Brief Description of the Drawings

## [0015]

5 Figure 1 depicts the changes of absorbance of the compound of the present invention (Compound a) and a compound disclosed in Japanese Patent Unexamined Publication No.(Hei) 7-70092/1995 (Compound b) when stored at 4°C.

Figure 2 depicts calibration curves of NADH obtained by absorption spectrum measurements. In the figures 1 and 2, ○ represents results obtained by using the compound of the present invention (Compound a), and □ represents results obtained by using the compound disclosed in Japanese Patent Unexamined Publication No.(Hei) 7-70092/1995 (Compound b).

Figure 3 depicts calibration curves of NADH obtained by absorption spectrum measurements. In the figure, ○ represents results obtained by using the compound of the present invention (Compound a), and □ represents results obtained by using nitro-TB.

15


## Best Mode for Carrying Out the Invention

[0016] As the alkyl group represented by R<sup>1</sup> a C<sub>1-4</sub> alkyl group is used. More specifically, methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group and other can be used. As the alkoxy group represented by R<sup>1</sup> a C<sub>1-4</sub> alkoxy group is used. More specifically, methoxy group, ethoxy group, n-propoxy group, isopropoxy group, n-butoxy group, sec-butoxy group, tert-butoxy group and other can be used. R<sup>1</sup> may preferably be methyl group or methoxy group.

[0017] As the alkali metal represented by M, for example, sodium, potassium and other can be used, and sodium is preferred.

25 [0018] The compounds of the present invention represented by the general formula (1) can be prepared by a conventional method. For example, a hydrazine compound represented by the following general formula (3):


30



35

can be reacted with an aldehyde compound represented by the general formula (4):


40



45


in an alcoholic solution to obtain a hydrazone compound represented by the general formula (5):

50



55

and then the resulting compound can be reacted with a corresponding diazonium salt in an organic solvent or water under basic condition to obtain a formazan compound represented by the general formula (6).



15 In the above general formula (6),

R<sup>1</sup> represents a C<sub>1-4</sub> alkyl group or a C<sub>1-4</sub> alkoxy group.

[0019] In the reactions mentioned above, sodium hydroxide, potassium hydroxide or other may be used as a basifying agent. Then, the resulting formazan compound of the general formula (6) can be oxidized by using an oxidizing agent such as butyl nitrite in an alcoholic solvent to obtain the tetrazolium salt compound of the general formula (1).

20 [0020] The compounds of the present invention have a characteristic feature that they receive a hydrogen produced by the action of a dehydrogenase via NADH or other, and they, per se, are reduced to form the formazan compounds represented by the above general formula (6). Accordingly, the concentration of a dehydrogenase such as lactate dehydrogenase, alcohol dehydrogenase, and glutamate dehydrogenase can be quantitatively measured by using the compounds of the present invention as hydrogen acceptors. They can also be used for quantitative measurement of

25 reducing substances such as NADH as described in the following examples. In general, the absorbance of the formazan compound of the general formula (6) formed in a reaction system may be measured to perform the quantitative measurement of these substances, or alternatively, the measurement can also be performed by means of fluorescent measurement or other spectrometric means. Methods for the measurement of a dehydrogenase utilizing an intermediate electronic transporter such as NADH are well known in the art, per se, and reaction conditions, measurement means and other can be appropriately chosen by one of ordinary skill in the art.

#### Examples

35 [0021] The present invention will be further explained more specifically by referring to the following examples. However, the scope of the present invention is not limited to the following examples.

Example 1: Synthesis of the compound of the general formula (1) wherein R<sup>1</sup> is methoxy group (Compound a)

40 [0022] p-Nitrophenylhydrazine (18.4 g) and sodium 4-formyl-1,3-benzenedisulfonate (37.2 g) were mixed with methanol, and the mixture was heated under reflux for two hours. The resulting precipitates were collected by filtration to obtain a hydrazone in a 85% yield. The hydrazone obtained (6.7 g) was dissolved in water (200 ml), and cooled to 0°C. Separately, 5-nitro-o-anisidine (2.7 g) was diazotized in a conventional manner, and the product was added to the hydrazone solution obtained above. The reaction mixture was kept at -5 to 0°C and added dropwise with an aqueous solution obtained by dissolving NaOH (2.6 g) in water (40 ml). After the dropwise addition, the mixture was stirred

45 overnight at room temperature. The reaction mixture was added with hydrochloric acid and concentrated. The precipitates formed by the addition of isopropanol were collected by filtration to obtain a formazan in a 63% yield. The resulting formazan (5 g) was suspended in methanol (250 ml), added with concentrated hydrochloric acid (6.7 ml) and butyl nitrite (4.1 g), and then the mixture was stirred at room temperature overnight. The reaction mixture was concentrated, and added with isopropanol. The deposited precipitates were collected by filtration, and the crude product was recrystallized from ethanol to obtain the captioned tetrazolium salt (Compound a) in a 45% yield.

50 Elemental analysis: Calculated. (C<sub>20</sub>H<sub>12</sub>N<sub>6</sub>O<sub>11</sub>S<sub>2</sub>Na) C: 40.00%, H: 2.18%, N: 14.00 %; Found. C: 39.78%, H: 2.15%, N: 13.76%

Example 2: Synthesis of the compound of the general formula (1) wherein R<sup>1</sup> is methyl group

55 [0023] The captioned compound was prepared in the same manner as in Example 1. Elemental analysis: Calculated. (C<sub>20</sub>H<sub>13</sub>N<sub>6</sub>O<sub>10</sub>S<sub>2</sub>Na) C: 41.10%, H: 2.24%, N: 14.38%; Found. C: 40.86%, H: 2.44%, N: 14.17%

## Example 3: Storage stability test

[0024] 50 mM Tris buffer (pH 8.0) containing 1 mM of Compound a was stored at 4°C for 0, 1, 7, 14, 31 and 90 days, and absorbances at 460 nm were measured. At the same time, the compound of the above general formula (2) described in Japanese Patent Unexamined Publication No. (Hei) 7-70092/1995 wherein both R<sup>1</sup> and R<sup>2</sup> are nitro groups (Compound b: see, the section of "Background Art") was stored in the same manner, and changes of absorbance were measured. Fig. 1 shows the relationships between the days of storage and absorbances deriving from Compound a and Compound b. Compound b exhibited remarkable increases of absorbance, whilst Compound a of the present invention gave no increase of absorbance for 90 days, which revealed its excellent stability.

## Example 4: Measurement of NADH concentration


[0025] 50 mM Tris buffer (pH 8.0, 5 ml) containing 0.1 mM of Compound a and 5 μM of 1-methoxy-5-methylphenazinium methosulfate was added with 0, 10, 20, 30, 40 or 50 μl of 5 mM NADH, and allowed to react at room temperature for 5 minutes, and then absorbances were measured. Figure 2 shows the relationship between NADH concentrations and absorbances. Measurement was performed in the same manner by using Compound b. Each of the compounds gave a linear calibration curve between NADH concentration and absorbance starting from the origin, and exhibited almost the same detection sensitivity. On the other hand, when the relationship between NADH concentration and absorbance was measured by using nitro-TB in the same manner, it was found that detection sensitivity was apparently inferior to that of the compound of the present invention as shown in Figure 3.

## Industrial Applicability

[0026] The tetrazolium salt compounds of the present invention give water-soluble formazans when used as hydrogen acceptors. Accordingly, no deposition is occurred in a measuring apparatus, and measurements by using automatic analyzers are facilitated. Furthermore, the compounds of the present invention have characteristics that they exhibit higher sensitivity compared to nitro-TB, which is generally used as a hydrogen acceptor in clinical diagnostics, and can easily detect NADH. In addition, they have the feature of excellent storage stability in the state of an aqueous solution. Accordingly, the compounds of the present invention are extremely useful as, for example, reagents for clinical diagnostics.

## Claims


1. Tetrazolium salt compound represented by the following general formula (1):



wherein R<sup>1</sup> is a C<sub>1-4</sub> alkyl group or a C<sub>1-4</sub> alkoxy group and M represents an alkali metal or an ammonium.

2. The tetrazolium salt compound according to claim 1 wherein R<sup>1</sup> is methyl group or methoxy group; and M is sodium.
3. A reagent for the measurement of a dehydrogenase which comprises a compound according to claim 1 or 2.
4. The reagent according to claim 3 which is used for a measuring process utilizing reduced nicotinamide adenine dinucleotide as an intermediate electron transporter.

5. A formazan compound represented by the following general formula (6):




wherein R<sup>1</sup> is a C<sub>1-4</sub> alkyl group or a C<sub>1-4</sub> alkoxy group.

6. A method for the measurement of a dehydrogenase wherein a compound according to claim 1 or 2 is used as a hydrogen acceptor.
7. The method according to claim 6 which comprises the step of measuring the absorbance of a formazan compound according to claim 5.
8. The method according to claim 6 or 7 wherein reduced nicotinamide adenine dinucleotide is used as an intermediate electronic transporter.


# Patentansprüche

1. Tetrazoliumsalz-Verbindung, dargestellt durch die folgende allgemeine Formel (1):



wobei R<sup>1</sup> eine C<sub>1-4</sub> Alkylgruppe oder eine C<sub>1-4</sub> Alkoxygruppe ist und M ein Alkalimetall oder ein Ammonium darstellt.

2. Die Tetrazoliumsalz-Verbindung nach Anspruch 1, wobei R<sup>1</sup> eine Methylgruppe oder Methoxygruppe ist; und M Natrium ist.
3. Ein Reagenz zur Messung einer Dehydrogenase, das die Verbindung gemäß Anspruch 1 oder 2 umfasst.
4. Das Reagenz nach Anspruch 3, das für ein Messverfahren verwendet wird unter Anwendung von reduziertem Nicotinamid-Adenin-Dinucleotid als intermediären Elektronenüberträger.
5. Eine Formazan-Verbindung, dargestellt durch die folgende allgemeine Formel (6):




(6)

wobei R<sup>1</sup> eine C<sub>1-4</sub> Alkylgruppe oder eine C<sub>1-4</sub> Alkoxygruppe ist.

6. Ein Verfahren zur Messung einer Dehydrogenase, **dadurch gekennzeichnet, dass** eine Verbindung gemäß Anspruch 1 oder 2 als Wasserstoffakzeptör verwendet wird.
7. Das Verfahren gemäß Anspruch 6, das den Schritt der Messung der Absorption einer Formazan-Verbindung gemäß Anspruch 5 umfasst.
8. Das Verfahren gemäß Anspruch 6 oder 7, **dadurch gekennzeichnet, dass** reduziertes Nicotinamid-Adenin-Dinucleotid als intermediärer Elektronenüberträger verwendet wird.

#### Revendications

1. Composé de sel de tétrazolium représenté par la formule générale suivante (1) :




(1)

dans laquelle R<sup>1</sup> est un groupe alkyle en C<sub>1-4</sub> ou un groupe alcoxy en C<sub>1-4</sub> et M représente un métal alcalin ou de l'ammonium.

2. Composé de sel de tétrazolium selon la revendication 1, dans lequel R<sup>1</sup> est un groupe méthyle ou un groupe méthoxy; et M est du sodium.
3. Réactif pour la mesure d'une déshydrogénase, qui comprend un composé selon la revendication 1 ou 2.
4. Réactif selon la revendication 3, qui est utilisé pour un procédé de mesure en utilisant un dinucléotide de nicotineamide adénine réduit comme transporteur d'électrons intermédiaire.
5. Composé de formazan représenté par la formule générale suivante (6) :





15 dans laquelle R<sup>1</sup> est un groupe alkyle en C<sub>1-4</sub> ou un groupe alcoxy en C<sub>1-4</sub>.

- 20 6. Procédé de mesure d'une déshydrogénase, dans lequel un composé selon la revendication 1 ou 2 est utilisé comme accepteur d'hydrogène.
- 25 7. Procédé selon la revendication 6, comprenant l'étape de mesure du pouvoir absorbant d'un composé de formazan selon la revendication 5.
8. Procédé selon la revendication 6 ou 7, dans lequel le dinucléotide de nicotinamide adénine réduit est utilisé comme transporteur d'électrons intermédiaire.




Fig. 2

